

## Mediquiz

### Series - 8

### The name of Cities in Medicine

1. Which of the following clinical conditions is diagnosed by the **PARIS** criteria?

- a. Autoimmune hepatitis-PBC overlap syndrome
- b. IgG4 associated disease
- c. Autoimmune pancreatitis
- d. Renal tubular acidosis

2. Which of the following is diagnosed with the help of **ROME** criteria?

- a. Inflammatory bowel disease
- b. Irritable bowel syndrome
- c. Premalignant lesions in the colon
- d. Hypoparathyroidism

3. Which disease is associated with the **WEST HAVEN** criteria?

- a. Sjogren syndrome
- b. Myocarditis
- c. Pleural mesothelioma
- d. Hepatic encephalopathy

4. Which clinical condition is associated with **NEW YORK** criteria ?

- a. Rheumatoid arthritis
- b. Renal vasculitis
- c. Psoriasis
- d. Ankylosing spondylitis

5. Which disease is associated with the **BERLIN** criteria?

- a. Hypopituitarism
- b. Scrub typhus
- c. ARDS
- d. Polychondritis

6. Recently, the "**OSLO DEFINITION**" has been formulated for which condition?

- a. SLE nephropathy
- b. Kaposi Sarcoma
- c. Celiac Disease
- d. TSH secreting tumour

7. Which public health topic was addressed in the **DOHA** declaration?

- a. Right to essential medicines
- b. Ethics in organ transplant
- c. Right to safe drinking water
- d. Palliative care in oncology

8. Which disease is diagnosed with the help of **MANCHESTER** criteria?

- a. Polyposis of colon
- b. Neurofibromatosis
- c. Autoimmune encephalitis
- d. Myocardial sarcoidosis



**Rudrajit Paul**  
Quiz Master

(Answer : next page)

**Answer : Mediquiz**

**Answers :-**

**1. A**

The Paris criteria is used for diagnosis of AIH-PBC overlap syndromes. These syndromes are rare and may be resistant to immune-suppressive therapy. The sensitivity and specificity of this criteria are both >90%.

**2. B**

The irritable bowel syndrome is diagnosed with the help of Rome criteria. The criteria were first published after a conference in Rome in 1988. But it has been modified over the years and new versions published. Currently, we are using the Rome IV criteria.

**3. D**

The West Haven criteria is used for diagnosis/ classification of hepatic encephalopathy. The classification system (also called modified Parsons-Smith classification) is given below:

Grade 0	No abnormality detected (mHE)
Grade 1	Trivial lack of awareness Euphoria or anxiety Shortened attention span Impairment of addition or subtraction
Grade 2	Lethargy Disorientation for time Obvious personality change Inappropriate behavior
Grade 3	Somnolence to semistupor Responsive to stimuli Confused
Grade 4	Gross disorientation, bizarre behavior Coma, unable to test mental state

(Source: Dove Medical Press)

**4. D**

Ankylosing spondylitis is diagnosed with the help of modified New York criteria. It has both clinical and radiological parameters

**5. C**

The Berlin criteria has been devised for ARDS in 2012. This criteria also includes classification of ARDS based on P/F ratio.

**6. C**

The Oslo definition has been formulated by a group of researchers for Celiac Disease. They have done extensive electronic database search for this purpose.

**7. A**

The Doha declaration is a landmark victory for public health activists of the world. By this agreement, a country is now able to grant licenses for drug production, in contravention of international intellectual property act, for public health emergencies.

**8. B**

The Manchester criteria was devised to diagnose NF type 2.

**There are only two sorts of doctors: those who practice with their brains, and those who practice with their tongues.**

**— William Osler: 1914 Quote**