

Mediquiz - 08 / 2021

Psychiatry

Dr Sujata Bhattacharya, MBBS, MD (Psy), Kolkata

(1) A patient sees a rope and gets afraid that it a snake. This sign is known as:

- a) Illusion
- b) Hallucination
- c) Delusion
- d) Depersonalization

(2) Schizophrenia of early onset which has bad prognosis is:

- a) Catatonic Schizophrenia
- b) Hebephrenic schizophrenia
- c) Paranoid schizophrenia
- d) Undifferentiated schizophrenia

(3) According to ICD for establishing a diagnosis of mania, the number of weeks the symptoms should persist for are:

- a) One
- b) Two
- c) Three
- d) Four

(4) The following drugs have abuse liability except:

- a) Buprenorphine
- b) Alprazolam
- c) Fluoxetine
- d) Dextropropoxyphene

(5) Which of the following is not suicide predictor?

- a) Adolescence
- b) Substance abuse
- c) Suicide note written
- d) Previous suicide attempt

(6) Treatment of choice of OCD is:

- a) Behaviour therapy
- b) Drug therapy
- c) Psychotherapy
- d) Combination of Behaviour and Drug therapy

(7) All are true about Conversion Disorder except:

- a) Presence of secondary gain
- b) Onset is late age
- c) Patient does not consciously produce symptoms
- d) Relation with stress

(8) AUDIT test is used for:

- a) Alcohol abuse disorder
- b) Opioid abuse disorder
- c) Sexual abuse disorder
- d) Cannabis use

(9) Which of the following the drug of choice for opioid overdose:

- a) Naloxone
- b) Clonidine
- c) Buprenorphine
- d) Flumazenil

(10) Reversible cause of dementia is:

- a) Hypothyroidism
- b) Alzheimer disease
- c) Vascular dementia
- d) Vitamin A deficiency

(11) Which of the following disorder is more common in female:

- a) Eating disorder
- b) ADHD
- c) Autism
- d) Conduct disorder

(12) Appetite for non-nutritive food is:

- a) Pica
- b) Anorexia
- c) Bulimia
- d) Binge

(13) Which of the following has least risk of causing movement disorder:

- a) Clozapine
- b) Olanzapine
- c) Quetiapine
- d) Paliperidone

(14) Which of the following hypnotic agents is least likely to be addictive:

- a) Ramelteon
- b) Zolpidem
- c) Temazepam
- d) Chlordiazepoxide

(15) How many days before surgery should Lithium be stopped:

- a) One
- b) Two
- c) Three
- d) Four

(Answer Page 77)